

Elastic Anomalies in Orbital-Degenerate Frustrated Spinel CoV_2O_4

Tadataka Watanabe^{1*}, Shogo Yamada¹, Rui Koborinai², Takuro Katsufuji²

¹*Department of Physics, College of Science and Technology, Nihon University, Chiyoda, Tokyo 101-8308, Japan*

²*Department of Physics, Waseda University, Shinjuku, Tokyo 169-8555, Japan*

Abstract

Ultrasound velocity measurements were performed on a single crystal of the orbital-degenerate frustrated spinel CoV_2O_4 in all the symmetrically-independent elastic moduli of the cubic crystal. The measurements of temperature dependence of the elastic moduli observed discontinuous elastic anomalies due to a ferrimagnetic transition at $T_C = 165$ K and another phase transition at $T^* = 50$ K. Additionally, the measurements observed anomalous temperature dependence of the elastic moduli, specifically, non-monotonic temperature dependence in the magnetically-ordered phase below T_C , and magnetic-field-sensitive elastic softening with decreasing temperature in the paramagnetic phase above T_C . These anomalous temperature variations below and above T_C should be driven by the coupling of lattice to magnetic excitations.

Keywords: Vanadate spinel, Frustration, Orbital order, Sound velocity

1 Introduction

Vanadate spinels AV_2O_4 ($A = \text{Zn}$ [1], Mg [2], Cd [3], Mn [4], Fe [5], and Co [6]) are geometrically frustrated magnets in which the pyrochlore lattice consists of V^{3+} ($3d^2$) ions with t_{2g} orbital degeneracy. AV_2O_4 undergoes a magnetic phase transition, specifically, an antiferromagnetic transition for nonmagnetic $A = \text{Zn}$ ($T_N = 40$ K [1]), Mg ($T_N = 42$ K [2]), and Cd ($T_N = 33$ K [3]), and a ferrimagnetic transition for magnetic $A = \text{Mn}$ ($T_C = 57$ K [4]), Fe ($T_C = 105$ K [5]), and Co ($T_C \sim 150$ K [6]). For AV_2O_4 , it is considered that the orbital degrees of freedom play an important role for the occurrence of the magnetic transition.

CoV_2O_4 is the only vanadate spinel which exhibits the absence of structural phase transition down to low temperature, while this compound exhibits a ferrimagnetic phase transition at $T_C \sim 150$ K [6]. And it is also noted that CoV_2O_4 exhibits the highest electrical conductivity among AV_2O_4 [6]. The

* Corresponding author. Tel: +81-3-3259-0920; fax: +81-3-3293-8269
E-mail address: tadataka@phys.cst.nihon-u.ac.jp

absence of structural phase transition and the nearly-metallic character in CoV₂O₄ are both expected to be results of the interplay between frustration and orbital degrees of freedom.

We performed ultrasound velocity measurements in CoV₂O₄ to study the interplay of orbital, spin, and lattice degrees of freedom. The sound velocity or the elastic modulus is a useful probe enabling symmetry-resolved thermodynamic information to be extracted from frustrated magnets [7,8,9,10,11]. The present study in CoV₂O₄ observed a variety of elastic anomalies which should have magnetic origins.

2 Experimental

The ultrasound velocity measurements were performed on a single crystal of CoV₂O₄ grown by the floating-zone method. Temperature (T) dependence of the ultrasound velocity was measured at T from 2 K to 300 K with magnetic field $H||[110]$ up to 7 T in all the symmetrically-independent elastic moduli of the cubic crystal: compression modulus C_{11} , tetragonal shear modulus $(C_{11}-C_{12})/2 = C_t$, and trigonal shear modulus C_{44} . The respective measurements of C_{11} , C_t , and C_{44} were performed using longitudinal sound waves with propagation $\mathbf{k}||[001]$ and polarization $\mathbf{u}||[001]$, transverse sound waves with $\mathbf{k}||[110]$ and $\mathbf{u}||[1-10]$, and transverse sound waves with $\mathbf{k}||[110]$ and $\mathbf{u}||[001]$. The sound velocities of CoV₂O₄ measured at room temperature (300 K) are 6740 m/s for C_{11} , 2750 m/s for C_t , and 3770 m/s for C_{44} .

3 Results and discussion

Figures 1(a), (b) and (c) respectively show T dependence of compression modulus $C_{11}(T)$, tetragonal shear modulus $C_t(T)$, and trigonal shear modulus $C_{44}(T)$ in CoV₂O₄ without magnetic field ($H = 0$). All the elastic modes exhibit a discontinuous change at $T_C = 165$ K as marked by arrows in Figs. 1(a)-(c), which is due to the ferrimagnetic transition. Additionally, only $C_{44}(T)$ exhibits a discontinuous change at $T^* = 50$ K as marked by arrow in Fig. 1(c), indicating the occurrence of another phase transition at T^* which sensitively couples to the trigonal lattice deformations. This




Figure 1: Temperature dependence of elastic moduli in CoV₂O₄ without magnetic field ($H = 0$). (a) Compression modulus C_{11} , (b) tetragonal shear modulus $(C_{11}-C_{12})/2 = C_t$, and (c) trigonal shear modulus C_{44} . T_C and T^* are marked by arrows in (a)-(c).

transition at T^* should correspond to the disorder-sensitive phase transition which was observed in the specific heat measurements [6,12].

As shown in Figs. 1(a)-(c), below T_C , all the elastic modes exhibit non-monotonic T dependence with characteristic minima at ~ 30 K and ~ 130 K for $C_{11}(T)$, ~ 30 K, ~ 100 K, and ~ 130 K for $C_t(T)$, and ~ 130 K for $C_{44}(T)$. Such an elastic-modulus-minimum anomaly in $C_t(T)$ usually arises from the coupling of lattice to gapped magnetic excitations [8,9,11]. Although the direct observation of magnetic excitations in CoV₂O₄ has not been reported yet, the elastic-modulus-minimum anomalies in $C_t(T)$ of CoV₂O₄ should stem from the coupling of lattice to gapped magnetic excitations. Since the temperature at which $C_t(T)$ minimum occurs is considered to correspond roughly to the gap magnitude of the excitations, the presence of the multiple $C_t(T)$ minima in the magnetically-ordered phase of CoV₂O₄ suggests the coupling of lattice to multiple gapped excitation modes.

In the paramagnetic phase ($T > T_C$), as shown in Figs. 1(a)-(c), $C_t(T)$ and $C_{44}(T)$ exhibit softening with decreasing T while $C_{11}(T)$ exhibits hardening with decreasing T . Figures 2(a), (b), and (c) show $C_{11}(T)$, $C_t(T)$, and $C_{44}(T)$ with $H \parallel [110]$ in the paramagnetic phase ($T > T_C$), respectively. It is evident that $C_t(T)$ varies with H in all the elastic modes. We note here that MnV₂O₄ and MgV₂O₄ exhibit H -sensitive elastic softening with decreasing T in the paramagnetic phase [10,11]. While $C_t(T)$ of MnV₂O₄ exhibits huge $\sim 1/T$ -type softening (convex T dependence) in the paramagnetic phase which is considered to be driven by orbital-spin-coupled fluctuations [10], $C_t(T)$ of MgV₂O₄ exhibits concave T dependence which is considered to be driven by orbital-spin-coupled gapped excitations [11]. Since the H -sensitive elastic softening of $C_t(T)$ and $C_{44}(T)$ in the paramagnetic phase of CoV₂O₄ resembles that of MgV₂O₄, this elastic softening in CoV₂O₄ should be attributed to orbital-spin-coupled gapped excitations. The H -sensitivity of $C_{11}(T)$ shown in Fig. 2(a) indicates that this elastic mode also contains the component of the H -sensitive softening which should be driven by the coupling of lattice to orbital-spin excitations, although the component of the H -sensitive softening in $C_{11}(T)$ is weaker than that of ordinal hardening (background) in this elastic mode. It is noted here that, as indicated by arrows in Figs. 2(a)-(c), while the softening in $C_t(T)$ and $C_{44}(T)$ is relaxed with increasing H , the softening component in $C_{11}(T)$ is enhanced with increasing H . Thus it is suggested that the softening component in $C_{11}(T)$ originates from orbital-spin-coupled excitations which are different from the origin of the softening in $C_t(T)$ and $C_{44}(T)$. Taking into account the absence of structural phase transition in CoV₂O₄ in contrast to its presence in MnV₂O₄ and MgV₂O₄, it is




Figure 2: Temperature dependence of elastic moduli in the paramagnetic phase of CoV₂O₄ ($T > T_C$) with $H \parallel [110]$. (a) $C_{11}(T)$, (b) $C_t(T)$, and (c) $C_{44}(T)$. The arrows in (a)-(c) are guides to the eye, indicating the variations of $C_{11}(T)$, $C_t(T)$, and $C_{44}(T)$ with increasing H .

expected that the orbital-spin-coupled excitations in CoV₂O₄ is inherently different from those in MnV₂O₄ and MgV₂O₄.

4 Summary

In summary, ultrasound velocity measurements in the orbital-degenerate frustrated spinel CoV₂O₄ observed a variety of elastic anomalies which should have magnetic origins. Discontinuous anomalies in T dependence of the elastic moduli at $T_C = 165$ K and $T^* = 50$ K should be due to a ferrimagnetic transition and another phase transition, respectively. Non-monotonic T dependence of the elastic moduli in the magnetically-ordered phase below T_C should be due to the coupling of lattice to gapped magnetic excitations. And H -sensitive elastic softening with decreasing T in the paramagnetic phase above T_C should be due to the coupling of lattice to orbital-spin-coupled excitations. Further experimental and theoretical studies are indispensable to understand the phase transitions and the magnetic excitations in CoV₂O₄.

5 Acknowledgement

This work was partly supported by a Grant-in-Aid for Scientific Research (C) (Grant No. 25400348) from MEXT of Japan, and by Nihon University College of Science and Technology Grants-in-Aid for Project Research.

References

- [1] S. H. Lee, D. Louca, H. Ueda, S. Park, T. J. Sato, M. Isobe, Y. Ueda, S. Rosenkranz, P. Zschack, J. Iniguez, Y. Qiu, R. Osborn, Phys. Rev. Lett. 93 (2004) 156407.
- [2] H. Mamiya, M. Onoda, T. Furubayashi, J. Tang, I. Nakatani, J. Appl. Phys. 81 (1997) 5289.
- [3] G. Giovannetti, A. Stroppa, S. Picozzi, D. Baldomir, V. Pardo, S. Blanco-Canosa, F. Rivadulla, S. Jodlauk, D. Niermann, J. Rohrkamp, T. Lorenz, S. Streltsov, D. I. Khomskii, J. Hemberger, Phys. Rev. B 83 (2011) 060402(R).
- [4] K. Adachi, T. Suzuki, K. Kato, K. Osaka, M. Takata, T. Katsufuji, Phys. Rev. Lett. 95 (2005) 197202.
- [5] T. Katsufuji, T. Suzuki, H. Takei, M. Shingu, K. Kato, K. Osaka, M. Takata, H. Sagayama, T. Arima, J. Phys. Soc. Jpn. 77 (2008) 053708.
- [6] A. Kismarhardja, J. S. Brooks, A. Kiswandhi, K. Matsubayashi, R. Yamanaka, Y. Uwatoko, J. Whalen, T. Siegrist, H. D. Zhou, Phys. Rev. Lett. 106 (2011) 056602.
- [7] T. Watanabe, S. Hara, and S. Ikeda, Phys. Rev. B 78 (2008) 094420.
- [8] T. Watanabe, S. Hara, S. Ikeda, and K. Tomiyasu, Phys. Rev. B 84 (2011) 020409(R).
- [9] T. Watanabe, S. Ishikawa, H. Suzuki, Y. Kousaka, and K. Tomiyasu, Phys. Rev. B 86 (2012) 144413.
- [10] Y. Nii, N. Abe, and T. Arima, Phys. Rev. B 87 (2013) 085111.
- [11] T. Watanabe, T. Ishikawa, S. Hara, A. T. M. N. Islam, E. M. Wheeler, and B. Lake, Phys. Rev. B 90 (2014) 100407(R).
- [12] Y. Huang, Z. Yang, and Y. Zhang, J. Phys: Condens. Matter 24 (2012) 056003.